

## SUMMARY OF PRODUCT CHARACTERISTICS

### 1. NAME OF THE MEDICINAL PRODUCT

DEMEPRAZOL 20 mg Capsules

### 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

**Active Substance:**

Omeprazole ..... 20 mg

**Excipients:**

Sucrose ..... 108.04 mg

For the full list of excipients, see section 6.1.

### 3. PHARMACEUTICAL FORM

Hard gelatin capsule

White-creamy enteric coated pellets in blue-colored hard gelatin capsules (No 1).

### 4. CLINICAL PARTICULARS

#### 4.1 Therapeutic indications

DEMEPRAZOL is indicated for:

Adults

- Treatment of duodenal ulcers
- Prevention of relapse of duodenal ulcers
- Treatment of gastric ulcers
- Prevention of relapse of gastric ulcers
- In combination with appropriate antibiotics, *Helicobacter pylori* (H. pylori) eradication in peptic ulcer disease
- Treatment of Non-steroidal anti-inflammatory drug (NSAID)-associated gastric and duodenal ulcers
- Prevention of NSAID-associated gastric and duodenal ulcers in patients at risk
- Treatment of reflux esophagitis
- Long-term management of patients with healed reflux esophagitis
- Treatment of symptomatic gastro-esophageal reflux disease (GERD)
- Treatment of Zollinger-Ellison syndrome

Pediatric use

*Children over 1 year of age and  $\geq 10$  kg*

- Treatment of reflux esophagitis
- Symptomatic treatment of heartburn and acid regurgitation in gastro-esophageal reflux disease

*Children and adolescents over 4 years of age*

- In combination with antibiotics in treatment of duodenal ulcer caused by *H. pylori*.

#### 4.2 Posology and method of administration

**Posology/frequency and duration of administration:**



### Posology in adults

#### *Treatment of duodenal ulcers*

The recommended dose in patients with an active duodenal ulcer is 20 mg once daily. In most patients healing occurs within two weeks. For those patients who may not be fully healed after the initial course, healing usually occurs during a further 2 weeks treatment period. In patients with poorly responsive duodenal ulcer DEMEPRAZOL 40 mg once daily is recommended and healing is usually achieved within 4 weeks.

#### *Prevention of relapse of duodenal ulcers*

For the prevention of relapse of duodenal ulcer in *H. pylori* negative patients or when *H. pylori* eradication is not possible the recommended dose is 20 mg once daily. In some patients a daily dose of 10 mg may be sufficient. In case of therapy failure, the dose can be increased to 40 mg.

#### *Treatment of gastric ulcers*

The recommended dose is 20 mg once daily. In most patients healing occurs within four weeks. For those patients who may not be fully healed after the initial course, healing usually occurs during a further 4 weeks treatment period. In patients with poorly responsive gastric ulcer DEMEPRAZOL 40 mg once daily is recommended and healing is usually achieved within 8 weeks.

#### *Prevention of relapse of gastric ulcers*

For the prevention of relapse in patients with poorly responsive gastric ulcer the recommended dose is 20 mg once daily. If needed, the dose can be increased to 40 mg once daily.

#### *H. pylori eradication in peptic ulcer disease*

For the eradication of *H. pylori* the selection of antibiotics should consider the individual patient's drug tolerance, and should be undertaken in accordance with national, regional and local resistance patterns and treatment guidelines.

- DEMEPRAZOL 20 mg + clarithromycin 500 mg + amoxicillin 1,000 mg, each twice daily for one week, or
- DEMEPRAZOL 20 mg + clarithromycin 250 mg (alternatively 500 mg) + metronidazole 400 mg (or 500 mg or tinidazole 500 mg), each twice daily for one week or
- DEMEPRAZOL 40 mg once daily with amoxicillin 500 mg and metronidazole 400 mg (or metronidazole 500 mg or tinidazole 500 mg), both three times a day for one week.

In each regimen, if the patient is still *H. pylori* positive, therapy may be repeated.

#### *Treatment of NSAID-associated gastric and duodenal ulcers*

For the treatment of NSAID-associated gastric and duodenal ulcers, the recommended dose is 20 mg once daily. In most patients healing occurs within 4 weeks. For those patients who may not be fully healed after the initial course, healing usually occurs during a further 4 weeks treatment period.

#### *Prevention of NSAID-associated gastric and duodenal ulcers in patients at risk*

For the prevention of NSAID-associated gastric ulcers or duodenal ulcers in patients at risk (age > 60, previous history of gastric and duodenal ulcers, previous history of upper GI bleeding) the recommended dose is 20 mg once daily.

#### *Treatment of reflux esophagitis*

The recommended dose is 20 mg once daily. In most patients healing occurs within four weeks. For those patients who may not be fully healed after the initial course, healing usually occurs during a



further 4 weeks treatment period.

In patients with severe esophagitis 40 mg once daily is recommended and healing is usually achieved within eight weeks.

*Long-term management of patients with healed reflux esophagitis*

For the long-term management of patients with healed reflux esophagitis the recommended dose is 10 mg once daily. If needed, the dose can be increased to 20-40 mg once daily.

*Treatment of symptomatic GERD*

The recommended dose is 20 mg daily. Patients may respond adequately to 10 mg daily, and therefore individual dose adjustment should be considered.

If symptom control has not been achieved after 4 weeks treatment with DEMEPRAZOL 20 mg daily, further investigation is recommended.

*Treatment of Zollinger-Ellison syndrome*

In patients with *Zollinger-Ellison* syndrome the dose should be individually adjusted and treatment continued as long as clinically indicated. The recommended initial dose is 60 mg daily. All patients with severe disease and inadequate response to other therapies have been effectively controlled and more than 90% of the patients maintained on doses of 20-120 mg daily. When dose exceed 80 mg daily, the dose should be divided and given twice daily.

**Route of Administration:**

It is recommended to take DEMEPRAZOL capsules in the morning, preferably without food, swallowed whole with half a glass of water. The capsules must not be chewed or crushed.

*For patients with swallowing difficulties and for children who can drink or swallow semi-solid food*

Patients can open the capsule and swallow the contents with half a glass of water or after mixing the contents in a slightly acidic fluid (e.g., fruit juice or apple juice, or in non-carbonated water) or in applesauce. Patients should be advised that the dispersion should be taken immediately (or within 30 minutes) and always be stirred just before drinking and rinsed down with half a glass of water.

Alternatively patients can suck the capsule and swallow the pellets with half a glass of water. The enteric-coated pellets must not be chewed.

**Additional information on special populations:**

**Renal impairment:**

Dose adjustment is not needed in patients with renal impairment (see section 5.2).

**Hepatic impairment:**

A daily dose of 10-20 mg may be sufficient in patients with hepatic impairment (see section 5.2).

**Pediatric population:**

*Children over 1 year of age and  $\geq 10$  kg*

*Treatment of reflux esophagitis*

*Symptomatic treatment of heartburn and acid regurgitation in gastro-esophageal reflux disease*

The posology recommendations are as follows:



Age	Weight	Posology
≥ 1 years	10-20 kg	10 mg once daily. The dose can be increased to 20 mg once daily if needed
≥ 2 years	> 20 kg	20 mg once daily. The dose can be increased to 40 mg once daily if needed

Reflux esophagitis: The treatment time is 4-8 weeks.

Symptomatic treatment of heartburn and acid regurgitation in GERD: The treatment time is 2–4 weeks. If symptom control has not been achieved after 2–4 weeks the patient should be investigated further.

#### Children and adolescents over 4 years of age

##### *Treatment of duodenal ulcer caused by H. pylori*

When selecting appropriate combination therapy, consideration should be given to official national, regional and local guidance regarding bacterial resistance, duration of treatment (most commonly 7 days but sometimes up to 14 days), and appropriate use of antibacterial agents.

The treatment should be supervised by a specialist.

The posology recommendations are as follows:

Weight	Posology
15–30 kg	Combination with two antibiotics: DEMEPRAZOL 10 mg, amoxicillin 25 mg/kg body weight and clarithromycin 7.5 mg/kg body weight are all administered together 2 times daily for 1 week
31–40 kg	Combination with two antibiotics: DEMEPRAZOL 20 mg, amoxicillin 750 mg and clarithromycin 7.5 mg/kg body weight are all administered 2 times daily for 1 week.
> 40 kg	Combination with two antibiotics: DEMEPRAZOL 20 mg, amoxicillin 1 g and clarithromycin 500 mg are all administered 2 times daily for 1 week.

#### **Geriatric population (>65 years)**

Dose adjustment is not needed in the elderly (see section 5.2).

#### **4.3 Contraindications**

Omeprazole is contraindicated in individuals hypersensitive to omeprazole, substituted benzimidazoles or to any of the excipients.

Omeprazole like other proton pump inhibitors (PPIs) must not be used concomitantly with nelfinavir (see section 4.5).

#### **4.4 Special warnings and precautions for use**

In the presence of any alarm symptom (e.g. significant unintentional weight loss, recurrent vomiting, dysphagia, hematemesis or melena) and when gastric ulcer is suspected or present, malignancy should be excluded, as treatment may alleviate symptoms and delay diagnosis.

Co-administration of atazanavir with proton pump inhibitors is not recommended (see section 4.5). If the combination of atazanavir with a proton pump inhibitor is judged unavoidable, close clinical monitoring (e.g. virus load) is recommended in combination with an increase in the dose of atazanavir to 400 mg with 100 mg of ritonavir; omeprazole 20 mg should not be exceeded.

Omeprazole, as all acid-blocking medicines, may reduce the absorption of vitamin B12 (cyanocobalamin) due to hypo- or achlorhydria. This should be considered in patients with reduced body stores or risk factors for reduced vitamin B12 absorption on long-term therapy.



Omeprazole is a CYP2C19 inhibitor. When starting or ending treatment with omeprazole, the potential for interactions with drugs metabolized through CYP2C19 should be considered. An interaction is observed between clopidogrel and omeprazole (see section 4.5). The clinical relevance of this interaction is uncertain. As a precaution, concomitant use of omeprazole and clopidogrel should be discouraged.

### **Hypomagnesemia**

Hypomagnesemia, symptomatic and asymptomatic, has been reported rarely in patients treated with PPIs for at least three months, in most cases after a year of therapy. Severe adverse events include tetany, arrhythmias and seizures. In most patients, treatment of hypomagnesemia required magnesium replacement and discontinuation of the PPI therapy. For patients expected to be on prolonged treatment or who take PPIs with digoxin or drugs that may cause hypomagnesaemia (e.g. diuretics), healthcare professionals should consider measuring magnesium levels before starting PPI treatment and periodically during treatment.

### **Bone fracture**

Observational studies suggest that proton pump inhibitor (PPI) therapy may increase the risk of osteoporosis-related fractures of hip, wrist and spine. The risk of fracture was increased in patients who received high-dose, defined as multiple doses, and long-term PPI therapy (a year or longer). Patients should use the lowest dose and shortest duration of PPI therapy appropriate to the conditions being treated.

Observational studies suggest that proton pump inhibitors may increase the overall risk of fracture by 10-40%. Some of this increase may be due to other risk factors. Patients at risk of osteoporosis should receive care according to current clinical guidelines and they should have an adequate intake of vitamin D and calcium.

### **Subacute cutaneous lupus erythematosus (SCLE)**

Proton pump inhibitors are associated with very infrequent cases of subacute cutaneous lupus erythematosus. If lesions occur, especially in sun-exposed areas of the skin, and if accompanied by arthralgia, the patient should seek medical help promptly and the health care professional should consider stopping DEMEPRAZOL treatments. Subacute cutaneous lupus erythematosus after previous treatment with a proton pump inhibitor may increase the risk of subacute cutaneous lupus erythematosus with other proton pump inhibitors.

### **Interactions with diagnostic investigations for neuroendocrine tumors**

Serum chromogranin A (CgA) levels increase secondary to drug-induced decreases in gastric acidity. The increased CgA level may cause false positive results in diagnostic investigations for neuroendocrine tumors. Healthcare providers should temporarily stop PPI treatment before assessing CgA levels and consider repeating the test if initial CgA levels are high. If serial tests are performed (e.g. for monitoring), the same commercial laboratory should be used for testing, as reference ranges between tests may vary.

Some children with chronic illnesses may require long-term treatment although it is not recommended.

Treatment with proton pump inhibitors may lead to slightly increased risk of gastrointestinal infections such as *Salmonella* and *Campylobacter* (see section 5.1).

As in all long-term treatments, especially when exceeding a treatment period of 1 year, patients should be kept under regular surveillance.

DEMEPRAZOL contains sucrose. Patients with rare hereditary problems of fructose intolerance, glucose/galactose malabsorption or sucrase-isomaltase insufficiency should not take this medicine.

#### **4.5 Interactions with other medicinal products and other forms of interaction**

##### Effects of omeprazole on the pharmacokinetics of other drugs

###### *Drugs with pH dependent absorption*

The decreased intragastric acidity during treatment with omeprazole might increase or decrease the absorption of drugs with a gastric pH dependent absorption.

###### *Nelfinavir, atazanavir*

The plasma levels of nelfinavir and atazanavir are decreased in case of co-administration with omeprazole. Concomitant administration of omeprazole with nelfinavir is contraindicated (see section 4.3). Co-administration of omeprazole (40 mg once daily) reduced mean nelfinavir exposure by ca. 40% and the mean exposure of the pharmacologically active metabolite M8 was reduced by ca. 75–90%. The interaction may also involve CYP2C19 inhibition.

Concomitant administration of omeprazole with atazanavir is not recommended (see section 4.4). Concomitant administration of omeprazole (40 mg once daily) and atazanavir 300 mg/ritonavir 100 mg to healthy volunteers resulted in a 75% decrease of the atazanavir exposure. Increasing the atazanavir dose to 400 mg did not compensate for the impact of omeprazole on atazanavir exposure. The co-administration of omeprazole (20 mg once daily) with atazanavir 400 mg/ritonavir 100 mg to healthy volunteers resulted in a decrease of approximately 30% in the atazanavir exposure as compared to atazanavir 300 mg/ritonavir 100 mg once daily.

###### *Digoxin*

Concomitant treatment with omeprazole (20 mg daily) and digoxin in healthy subjects increased the bioavailability of digoxin by 10%. Digoxin toxicity has been rarely reported. However caution should be exercised when omeprazole is given at high doses in elderly patients. Therapeutic drug monitoring of digoxin should be then reinforced.

###### *Clopidogrel*

Results from studies in healthy subjects have shown a pharmacokinetic/pharmacodynamic interaction between clopidogrel (300 mg loading dose/75 mg daily maintenance dose) and omeprazole (80 mg p.o. daily) resulting in a decreased exposure to the active metabolite of clopidogrel by an average of 46% and a decreased maximum inhibition of (ADP induced) platelet aggregation by an average of 16%.

Inconsistent data on the clinical implications of a pharmacokinetic/pharmacodynamic interaction of omeprazole in terms of major cardiovascular events have been reported from both observational and clinical studies. As a precaution, concomitant use of omeprazole and clopidogrel should be discouraged (see section 4.4).

###### *Other Drugs*

The absorption of posaconazole, erlotinib, ketoconazole and itraconazole is significantly reduced and thus clinical efficacy may be impaired. Concomitant use of posaconazole and erlotinib should be avoided.



### Drugs metabolized by CYP2C19

Omeprazole is a moderate inhibitor of CYP2C19, the major omeprazole metabolizing enzyme. Thus, the metabolism of concomitant active substances also metabolized by CYP2C19, may be decreased and the systemic exposure to these substances increased. Examples of such drugs are R-warfarin and other vitamin K antagonists, cilostazol, diazepam and phenytoin.

#### *Cilostazol*

Omeprazole, given in doses of 40 mg to healthy subjects in a cross-over study, increased  $C_{max}$  and area under the plasma concentration-time curve (AUC) levels for cilostazol by 18% and 26% respectively, and  $C_{max}$  and AUC levels of one of its active metabolites by 29% and 69% respectively.

#### *Phenytoin*

Monitoring phenytoin plasma concentration is recommended during the first two weeks after initiating omeprazole treatment and, if a phenytoin dose adjustment is made, omeprazole treatment should be monitored and a further dose adjustment should be made.

### Unknown mechanism

#### *Saquinavir*

Concomitant administration of omeprazole with saquinavir/ritonavir resulted in increased plasma levels up to approximately 70% for saquinavir associated with good tolerability in HIV-infected patients.

#### *Tacrolimus*

Concomitant administration of omeprazole has been reported to increase the serum levels of tacrolimus. A reinforced monitoring of tacrolimus concentrations as well as renal function (creatinine clearance) should be performed, and dosage of tacrolimus adjusted if needed.

#### *Methotrexate*

When given together with PPIs, methotrexate levels have been reported to increase in some patients. In high-dose methotrexate administration a temporary withdrawal of omeprazole may need to be considered.

### Effects of other drugs on the pharmacokinetics of omeprazole

#### *Inhibitors CYP2C19 and/or CYP3A4*

Since omeprazole is metabolized by CYP2C19 and CYP3A4, active substances known to inhibit CYP2C19 or CYP3A4 (such as clarithromycin and voriconazole) may lead to increased omeprazole serum levels by decreasing omeprazole's rate of metabolism. Concomitant voriconazole treatment resulted in more than doubling of the omeprazole exposure. As high doses of omeprazole have been well-tolerated adjustment of the omeprazole dose is not generally required. However, dose adjustment should be considered in patients with severe hepatic impairment and if long-term treatment is indicated.

#### *Inducers of CYP2C19 and/or CYP3A4*

Drugs known to induce CYP2C19 or CYP3A4 or both (such as rifampicin and St John's Wort) may lead to decreased omeprazole serum levels by increasing omeprazole's rate of metabolism.

### **Information regarding special populations**



### **Pediatric population**

There is no sufficient information to recommend omeprazole to pediatric populations.

## **4.6 Fertility, pregnancy and lactation**

### **General Recommendation**

Pregnancy category is C.

### **Women with child-bearing potential/Contraception**

No special advice is given for use of this drug in women with child-bearing potential and practicing contraception.

### **Pregnancy**

Results from three prospective epidemiological studies (more than 1000 exposed outcomes) indicate no adverse effects of omeprazole on pregnancy or on the health of the fetus/newborn child. Animal studies are insufficient with regard to effects on pregnancy and/or embryonal/fetal development and/or parturition and/or post-natal development (see section 5.3). Potential risk to human is unknown.

DEMEPRAZOL should only be used during pregnancy if the benefits to the mother outweigh the potential risks to the fetus.

### **Breast-feeding**

Omeprazole is excreted in breast milk; however it is not likely to influence the child when therapeutic doses are used.

### **Fertility**

Animal studies with the racemic mixture omeprazole, given by oral administration do not indicate effects with respect to fertility.

## **4.7 Effects on ability to drive and use machines**

DEMEPRAZOL is not likely to affect the ability to drive or use machines. Adverse drug reactions such as dizziness and visual disturbances may occur (see section 4.8). If affected, patients should not drive or operate machinery.

## **4.8 Undesirable effects**

The most common side effects (1-10% of patients) are headache, abdominal pain, constipation, diarrhea, flatulence and nausea/vomiting.

The following adverse drug reactions have been identified or suspected in the clinical trials program for omeprazole and post-marketing. None was found to be dose-related. Adverse reactions listed below are classified according to frequency and System Organ Class (SOC).

Frequency categories are defined according to the following convention:

Very common ( $\geq 1/10$ ), Common ( $\geq 1/100$  to  $< 1/10$ ), Uncommon ( $\geq 1/1,000$  to  $< 1/100$ ), Rare ( $\geq 1/10,000$  to  $< 1/1,000$ ), Very rare ( $< 1/10,000$ ), Not known (cannot be estimated from the available data).

### **Blood and lymphatic system disorders**

*Rare:* Leucopenia, thrombocytopenia

*Very rare:* Agranulocytosis, pancytopenia



### **Immune system disorders**

*Rare:* Hypersensitivity reactions e.g. fever, angioedema and anaphylactic reaction/shock

### **Metabolism and nutrition disorders**

*Rare:* Hyponatremia

*Not known:* Hypomagnesemia; severe hypomagnesemia may result in hypocalcemia. Hypomagnesemia may also be associated with hypokalemia.

### **Psychiatric disorders**

*Uncommon:* Insomnia

*Rare:* Agitation, confusion, depression, indisposition.

*Very rare:* Aggression, hallucinations, tiredness

### **Nervous system disorders**

*Common:* Headache

*Uncommon:* Dizziness, paresthesia, somnolence

*Rare:* Taste disturbance

*Not known:* Reversible orientation disorder

### **Eye disorders**

*Rare:* Blurred vision

*Very rare:* Irreversible visual impairment (observed in isolated cases in critically patients receiving intravenous omeprazole injection, causal relationship has not been established)

### **Ear and labyrinth disorders**

*Uncommon:* Vertigo

### **Cardiac disorders**

*Very rare:* Chest pain (may be similar to angina pectoris)

### **Respiratory, thoracic and mediastinal disorders**

*Rare:* Bronchospasm

### **Gastrointestinal disorders**

*Common:* Abdominal pain, constipation, diarrhea, flatulence, nausea/vomiting, fundic gland polyps (benign)

*Rare:* Dry mouth, stomatitis, gastrointestinal candidiasis

*Very rare:* Polychromosomal hypertrophy of the gastric mucosal membrane, most cases have been defined as benign and reversible gland-like cysts.

*Not known:* Microscopic colitis, serum gastrin levels increased in the first 2 weeks of omeprazole treatment, but returned to normal after end of treatment.

### **Hepatobiliary disorders**

*Uncommon:* Increased liver enzymes

*Rare:* Hepatitis with or without jaundice

*Very rare:* Hepatic failure, encephalopathy in patients with pre-existing liver disease

### **Skin and subcutaneous tissue disorders**

*Uncommon:* Dermatitis, pruritus, rash, urticaria

*Rare:* Alopecia, photosensitivity



*Very rare:* Erythema multiforme, Stevens-Johnson syndrome, toxic epidermal necrolysis (TEN)  
*Unknown:* Subacute cutaneous lupus erythematosus (see section 4.4)

#### **Musculoskeletal and connective tissue disorders**

*Uncommon:* Fracture of the hip, wrist or spine  
*Rare:* Arthralgia, myalgia  
*Very rare:* Muscular weakness, rhabdomyolysis

#### **Renal and urinary disorders**

*Rare:* Interstitial nephritis

#### **Reproductive system and breast disorders**

*Very rare:* Gynecomasty  
*Unknown:* Sexual function disorders

#### **General disorders and administration site conditions**

*Uncommon:* Malaise, peripheral edema  
*Rare:* Hyperhidrosis

#### **Additional information on special populations**

##### **Pediatric population**

The safety of omeprazole has been assessed in a total of 310 children aged 0 to 16 years with acid-related disease. There are limited long term safety data from 46 children who received maintenance therapy of omeprazole during a clinical study for severe erosive esophagitis for up to 749 days. The adverse event profile was generally the same as for adults in short- as well as in long-term treatment. There are no long term data regarding the effects of omeprazole treatment on puberty and growth.

##### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorization of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions in accordance with local requirements.

#### **4.9 Overdose**

There is limited information available on the effects of overdoses of omeprazole in humans. In the literature, doses of up to 560 mg have been described, and occasional reports have been received when single oral doses have reached up to 2,400 mg omeprazole (120 times the usual recommended clinical dose). Nausea, vomiting, dizziness, abdominal pain, diarrhea and headache have been reported. Also apathy, depression and confusion have been described in single cases.

The symptoms described have been transient, and no serious outcome has been reported. The rate of elimination was unchanged (first order kinetics) with increased doses. Treatment, if needed, is symptomatic.

### **5. PHARMACOLOGICAL PROPERTIES**

#### **5.1 Pharmacodynamic properties**

**Pharmacotherapeutic group:** Proton pump inhibitors



**ATC code:** A02BC01

#### Mechanism of action

Omeprazole, a racemic mixture of two enantiomers reduces gastric acid secretion through a highly targeted mechanism of action. It is a specific inhibitor of the acid pump in the parietal cell. It is rapidly acting and provides control through reversible inhibition of gastric acid secretion with once daily dosing.

Omeprazole is a weak base and is concentrated and converted to the active form in the highly acidic environment of the intracellular canaliculi within the parietal cell, where it inhibits the enzyme H<sup>+</sup>K<sup>+</sup>-ATPase (the acid pump). This effect on the final step of the gastric acid formation process is dose-dependent and provides for highly effective inhibition of both basal acid secretion and stimulated acid secretion, irrespective of stimulus.

#### Pharmacodynamic effects

All pharmacodynamic effects observed can be explained by the effect of omeprazole on acid secretion.

#### *Effect on gastric acid secretion*

Oral dosing with omeprazole once daily provides for rapid and effective inhibition of daytime and night-time gastric acid secretion with maximum effect being achieved within 4 days of treatment. With omeprazole 20 mg, a mean decrease of at least 80% in 24-hour intragastric acidity is then maintained in duodenal ulcer patients, with the mean decrease in peak acid output after pentagastrin stimulation being about 70% 24 hours after dosing.

Oral dosing with omeprazole 20 mg maintains an intragastric pH of  $\geq 3$  for a mean time of 17 hours of the 24-hour period in duodenal ulcer patients.

As a consequence of reduced acid secretion and intragastric acidity, omeprazole dose-dependently reduces/normalizes acid exposure of the esophagus in patients with GERD.

The inhibition of acid secretion is related to the AUC of omeprazole and not to the actual plasma concentration at a given time.

No tachyphylaxis has been observed during treatment with omeprazole.

#### *Effect on H. pylori*

*H. pylori* is associated with peptic ulcer disease, including duodenal and gastric ulcer disease. *H. pylori* is a major factor in the development of gastritis. *H. pylori* together with gastric acid are major factors in the development of peptic ulcer disease. *H. pylori* is a major factor in the development of atrophic gastritis which is associated with an increased risk of developing gastric cancer.

Eradication of *H. pylori* with omeprazole and antimicrobials is associated with, high rates of healing and long-term remission of peptic ulcers.

Dual therapies have been tested and found to be less effective than triple therapies. They could, however, be considered in cases where known hypersensitivity precludes use of any triple combination.

#### *Other effects related to acid inhibition*



During long-term treatment gastric glandular cysts have been reported in a somewhat increased frequency. These changes are a physiological consequence of pronounced inhibition of acid secretion, are benign and appear to be reversible.

Decreased gastric acidity due to any means including proton pump inhibitors, increases gastric counts of bacteria normally present in the gastrointestinal tract. Treatment with acid-reducing drugs may lead to slightly increased risk of gastrointestinal infections such as *Salmonella* and *Campylobacter* and possibly *Clostridium difficile* in in-patients.

During treatment with medicinal products inhibiting acid secretion, serum gastrin levels increases in response to the decreased acid secretion. Also CgA increases due to decreased gastric acidity. The increased CgA level may interfere with investigations for neuroendocrine tumours. Available published evidence suggests that proton pump inhibitors should be discontinued between 5 days and 2 weeks prior to CgA measurement. This is to allow CgA levels that might be spuriously elevated following PPI treatment to return to reference range.

An increased number of enterochromaffin-like (ECL) cells possibly related to the increased serum gastrin levels, have been observed in some patients (both children and adults) during long term treatment with omeprazole. The findings are considered to be of no clinical significance.

#### Pediatric use

In a non-controlled study in children (1 to 16 years of age) with severe reflux esophagitis, omeprazole at doses of 0.7 to 1.4 mg/kg improved esophagitis level in 90% of the cases and significantly reduced reflux symptoms. In a single-blind study, children aged 0–24 months with clinically diagnosed GERD were treated with 0.5, 1.0 or 1.5 mg omeprazole/kg. The frequency of vomiting/regurgitation episodes decreased by 50% after 8 weeks of treatment irrespective of the dose.

#### *Eradication of *H. pylori* in children*

A randomized, double blind clinical study (Héliot study) concluded that omeprazole in combination with two antibiotics (amoxicillin and clarithromycin), was safe and effective in the treatment of *H. pylori* infection in children age 4 years old and above with gastritis: *H. pylori* eradication rate: 74.2% (23/31 patients) with omeprazole + amoxicillin + clarithromycin versus 9.4% (3/32 patients) with amoxicillin + clarithromycin. However, there was no evidence of any clinical benefit with respect to dyspeptic symptoms. This study does not support any information for children aged less than 4 years.

## **5.2 Pharmacokinetic properties**

### **General properties**

#### Absorption:

Omeprazole and omeprazole magnesium are acid labile and are therefore administered orally as enteric-coated granules in capsules or tablets. Absorption of omeprazole is rapid, with peak plasma levels occurring approximately 1-2 hours after dose. Absorption of omeprazole takes place in the small intestine and is usually completed within 3-6 hours. Concomitant intake of food has no influence on the bioavailability. The systemic availability (bioavailability) from a single oral dose of omeprazole is approximately 40%. After repeated once-daily administration, the bioavailability increases to about 60%.



Distribution:

The apparent volume of distribution in healthy subjects is approximately 0.3 L/kg body weight. Omeprazole is 97% plasma protein bound.

Biotransformation:

Omeprazole is completely metabolized by the cytochrome P450 system (CYP). The major part of its metabolism is dependent on the polymorphically expressed CYP2C19, responsible for the formation of hydroxyomeprazole, the major metabolite in plasma. The remaining part is dependent on another specific isoform, CYP3A4, responsible for the formation of omeprazole sulphone. As a consequence of high affinity of omeprazole to CYP2C19, there is a potential for competitive inhibition and metabolic drug-drug interactions with other substrates for CYP2C19. However, due to low affinity to CYP3A4, omeprazole has no potential to inhibit the metabolism of other CYP3A4 substrates. In addition, omeprazole lacks an inhibitory effect on the main CYP enzymes.

Approximately 3% of the Caucasian population and 15-20% of Asian populations lack a functional CYP2C19 enzyme and are called poor metabolizers. In such individuals the metabolism of omeprazole is probably mainly catalyzed by CYP3A4. After repeated once-daily administration of 20 mg omeprazole, the mean AUC was 5 to 10 times higher in poor metabolizers than in subjects having a functional CYP2C19 enzyme (extensive metabolizers). Mean peak plasma concentrations were also higher, by 3 to 5 times. These findings have no implications for the posology of omeprazole.

Elimination:

The plasma elimination half-life of omeprazole is usually shorter than one hour both after single and repeated oral once-daily dosing. Omeprazole is completely eliminated from plasma between doses with no tendency for accumulation during once-daily administration. Almost 80% of an oral dose of omeprazole is excreted as metabolites in the urine, the remainder in the feces, primarily originating from bile secretion.

Linearity/Non-linearity

The AUC of omeprazole increases with repeated administration. This increase is dose-dependent and results in a non-linear dose-AUC relationship after repeated administration. This time- and dose-dependency is due to a decrease of first pass metabolism and systemic clearance probably caused by an inhibition of the CYP2C19 enzyme by omeprazole and/or its metabolites (e.g. the sulphone).

No metabolite has been found to have any effect on gastric acid secretion.

**Characteristics in patients**

Hepatic impairment:

The metabolism of omeprazole in patients with liver dysfunction is impaired, resulting in an increased AUC. Omeprazole has not shown any tendency to accumulate with once daily dosing.

Renal impairment:

The pharmacokinetics of omeprazole, including systemic bioavailability and elimination rate, are unchanged in patients with reduced renal function.

Elderly:

The metabolism rate of omeprazole is somewhat reduced in elderly subjects (75-79 years of age).

Pediatric patients:

During treatment with the recommended doses to children from the age of 1 year, similar plasma concentrations were obtained as compared to adults. In children younger than 6 months, clearance of omeprazole is low due to low capacity to metabolize omeprazole.

### **5.3 Preclinical safety data**

Gastric ECL-cell hyperplasia and carcinoids, have been observed in life-long studies in rats treated with omeprazole. These changes are the result of sustained hypergastrinaemia secondary to acid inhibition. Similar findings have been made after treatment with H<sub>2</sub>-receptor antagonists, proton pump inhibitors and after partial fundectomy. Thus, these changes are not from a direct effect of any individual medicine.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

#### Sugar spheres:

- Sucrose
- Maize starch
- Purified Water

Sodium lauryl sulphate

Anhydrous disodium phosphate

Mannitol (E421)

Hypromellose (E464)

Macrogol 6000

Talc (E553b)

Polysorbate 80 (E433)

Titanium dioxide (E171)

Eudragit L30-D55

#### Hard gelatin capsule (Bovine bone gelatin)

Titanium dioxide

Yellow iron oxide

Indigotine FD&C blue 2

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

36 months.

### **6.4 Special precautions for storage**

Store at room temperature 25°C in a dry place. Capsules must be used within 1 month after package is opened; if any medicine remains when this time period is exceeded, do not use.

### **6.5 Nature and contents of container**

30 ml white opaque HDPE bottle with white opaque polypropylene child-resistant screw-cap, which contains 14 capsules.

### **6.6 Special precautions for disposal and other handling**

Any unused material should be disposed according to local disposal regulations.



**7. MARKETING AUTHORIZATION HOLDER**

DEVA Holding A.Ş.  
Halkalı Merkez Mah. Basın Ekspres Cad. No:1  
34303 Küçükçekmece-ISTANBUL/TÜRKİYE

**8. MARKETING AUTHORIZATION NUMBER(S)**

155/57

**9. DATE OF FIRST AUTHORIZATION/RENEWAL OF THE AUTHORIZATION**

Date of first authorization : 22.03.1991  
Date of latest renewal : 20.03.2013

**10. DATE OF REVISION OF THE TEXT**